

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/339049527>

An Integrated Care Intervention Supported by a Mobile Health Tool in Patients Using Noninvasive Ventilation at Home: Randomized Controlled Trial (Preprint)

Article in *JMIR mhealth and uhealth* · September 2019

DOI: 10.2196/16395

CITATIONS

4

READS

85

12 authors, including:



Erik Baltaxe

Rabin Medical Center

30 PUBLICATIONS 202 CITATIONS

[SEE PROFILE](#)



Eva Aumatell

Universitat Oberta de Catalunya

7 PUBLICATIONS 27 CITATIONS

[SEE PROFILE](#)



Anael Barberan-Garcia

77 PUBLICATIONS 837 CITATIONS

[SEE PROFILE](#)



Carmen Herranz

Consorci d'Atenció Primària de Salut-Barcelona Esquerra CAPSBE

4 PUBLICATIONS 23 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Regional Implementation of Collaborative Lung Function Testing [View project](#)



Synergy-COPD [View project](#)

Original Paper

Integrated Care Intervention Supported by a Mobile Health Tool for Patients Using Noninvasive Ventilation at Home: Randomized Controlled Trial

Erik Baltaxe^{1,2}, MD; Cristina Embid^{1,2}, MD, PhD; Eva Aumatell¹, MSc; María Martínez¹, MSc; Anael Barberan-García^{1,2}, PhD; John Kelly³, MSc; John Eaglesham³, MSc; Carmen Herranz^{1,2}, MSc; Eloisa Vargiu⁴, PhD; Josep Maria Montserrat^{1,2}, MD, PhD; Josep Roca^{1,2}, MD, PhD; Isaac Cano^{1,2}, PhD

¹Hospital Clínic de Barcelona, Institut d'Investigacions Biomèdiques August Pi i Sunyer, Universitat de Barcelona, Barcelona, Spain

²Center for Biomedical Research Network in Respiratory Diseases, Madrid, Spain

³Advanced Digital Innovation (UK) Ltd, Salts Mill, United Kingdom

⁴Eurecat Technological Center of Catalonia, Barcelona, Spain

Corresponding Author:

Isaac Cano, PhD

Hospital Clínic de Barcelona

Institut d'Investigacions Biomèdiques August Pi i Sunyer

Universitat de Barcelona

Villarroel 170

Barcelona, 08036

Spain

Phone: 34 932275540

Email: ISCANO@clinic.cat

Abstract

Background: Home-based noninvasive ventilation has proven cost-effective. But, adherence to therapy still constitutes a common clinical problem. We hypothesized that a behavioral intervention supported by a mobile health (mHealth) app could enhance patient self-efficacy. It is widely accepted that mHealth-supported services can enhance productive interactions among the stakeholders involved in home-based respiratory therapies.

Objective: This study aimed to measure changes in self-efficacy in patients with chronic respiratory failure due to diverse etiologies during a 3-month follow-up period after the intervention. Ancillary objectives were assessment of usability and acceptability of the mobile app as well as its potential contribution to collaborative work among stakeholders.

Methods: A single-blind, single-center, randomized controlled trial was conducted between February 2019 and June 2019 with 67 adult patients with chronic respiratory failure undergoing home-based noninvasive ventilation. In the intervention group, a psychologist delivered a face-to-face motivational intervention. Follow-up was supported by a mobile app that allowed patients to report the number of hours of daily noninvasive ventilation use and problems with the therapy. Advice was automatically delivered by the mobile app in case of a reported problem. The control group received usual care. The primary outcome was the change in the Self Efficacy in Sleep Apnea questionnaire score. Secondary outcomes included app usability, app acceptability, continuity of care, person-centered care, and ventilatory parameters.

Results: Self-efficacy was not significantly different in the intervention group after the intervention (before: mean 3.4, SD 0.6; after: mean 3.4, SD 0.5, $P=.51$). No changes were observed in adherence to therapy nor quality of life. Overall, the mHealth tool had a good usability score (mean 78 points) and high acceptance rate (mean score of 7.5/10 on a Likert scale). It was considered user-friendly (mean score of 8.2/10 on a Likert scale) and easy to use without assistance (mean score of 8.5/10 on a Likert scale). Patients also scored the perception of continuity of care and person-centered care as high.

Conclusions: The integrated care intervention supported by the mobile app did not improve patient self-management. However, the high acceptance of the mobile app might indicate potential for enhanced communication among stakeholders. The study identified key elements required for mHealth tools to provide effective support to collaborative work and personalized care.

Trial Registration: ClinicalTrials.gov NCT03932175; <https://clinicaltrials.gov/ct2/show/NCT03932175>

KEYWORDS

behavioral change; eHealth; noninvasive ventilation; mobile health; chronic diseases

Introduction

In the 1950s, the polio epidemic demonstrated the safety and efficacy of noninvasive ventilation (NIV) to decrease mortality [1]. Since then, the use of this therapeutic approach at home has reduced hospital admissions, has favorably impacted health-related quality of life, improved sleep quality, and reduced mortality in patients with chronic respiratory failure due to diverse etiologies [2-8]. These results have driven a steady increase in the prevalence of patients using home-based NIV in Europe, ranging from 4.5 to 20 per 100,000 adults [9-11].

Despite its proven cost-effectiveness [12], patient adherence to home-based NIV could still improve, which should further enhance health care-related efficiencies of the intervention [13]. Monitoring and optimization of physiological settings have enhanced adherence by improving the timely detection of problems such as mask leaks and patient-ventilator asynchronies [14]. Nevertheless, improvement in the behavioral aspects such as patient motivation and empowerment for self-management are important factors to consider when addressing adherence to respiratory therapies.

The current study sought to explore the transfer of previous positive experiences with behavioral interventions in other fields (ie, physical activity) [15-20] to home-based NIV. Specifically, we addressed the concept of self-efficacy, defined as the individual's perceived capability to perform a particular behavior [21]. Self-efficacy expectations can be affected by enablers or barriers such as the perception of physical function or the capacity for self-management. Therefore, a person who does not believe in her or his capacity to perform the desired action will fail to adopt, initiate, and maintain it. Self-efficacy is therefore seen as the most influential motivational factor and the strongest predictor of behavioral intentions [21].

We propose the use of a behavioral mobile health (mHealth) intervention, which can be framed by Bandura's model [22], to support changes in self-efficacy. This model is based on the concepts of health risk perceptions, health outcome expectancies, and the patients' confidence to engage in certain behaviors. The model has been widely applied in studies of the adoption, initiation, and maintenance of health-promoting behaviors [23].

In addition to self-efficacy as a way to influence behavioral change, previous reports by Hernandez et al [24] and Cano et al [25] identified two common hinderances for the effective implementation of complex respiratory therapies (ie, long-term oxygen therapy, continuous positive airway pressure therapy, home NIV, and home-based nebulizer therapy). First, interaction and communication, which could greatly benefit from digital tools supporting collaborative work, are needed among several stakeholders, namely health professionals at different health care tiers (eg, primary care, specialized care), patients and carers, companies undertaking equipment maintenance, and others. Second, improvement in therapeutic adherence is needed, which

could be achieved by empowering patients to perform self-management.

Within this context, information and communication technologies (ICT) have been identified as promising tools to enhance the coordination between stakeholders and contribute to improved health outcomes [26,27]. Nonetheless, the implementation remains immature [28] due to a lack of evidence in a real-world context for the capacity of ICT to sustain behavioral changes, including self-efficacy, in patients with chronic, complex conditions. It is widely accepted that, despite current limitations, patients with chronic, complex conditions are an ideal population for which care coordination, patient and medical staff satisfaction, and patient empowerment are of the utmost importance to produce health benefits.

The principal objective of this study was to explore the capacity of a behavioral mHealth intervention to increase patient empowerment for self-management and adherence to therapy. The secondary aim was to learn, based on the experience of professionals and patients, how the mHealth tool should evolve to support collaborative work.

Methods

Study Design and Participants

A single-blind, single-center, randomized controlled trial with two parallel arms (1:1 ratio) was conducted. Patients were randomized to a control group or an intervention arm, which consisted of the behavioral mHealth intervention in addition to usual care. Inclusion criteria were as follows: all adult patients with hypercapnic ventilatory failure due to chest wall, neuromuscular, lung parenchyma, or airway disease already receiving treatment with NIV irrespective of treatment duration and in possession of a mobile phone or tablet that could support the use of the mHealth app (MyPathway). MyPathway [29] is a secure, digital communication channel connecting patients to clinicians and services. It is an app-based tool for both patients and clinicians to use on phones or tablets. See [Multimedia Appendix 1](#) for more details. Patients with severe psychiatric or neurological diseases were excluded, as well as patients hospitalized at the time of assessment.

Intervention

In addition to usual care, the behavioral mHealth intervention included a face-to-face motivational interview by a psychologist (EA) to assess the patient's adherence profile and lifestyle habits, with a follow-up through the MyPathway app. In contrast, the control group received only usual care, which consisted of manual discharge and review of the NIV machine data by the treating pulmonologist and respiratory nurse. Respiratory parameters were changed, if needed, according to clinical data (anamnesis and physical examination) in addition to NIV data.

At the time of enrollment, semi-structured motivational interviews were conducted individually. Participants were asked about their treatment adaptation experience, lifestyle (physical activity and food habits), and use of ICT. In each session, field notes were taken anonymously, and no recordings were made. The intervention consisted of a 10-50-minute face-to-face session at the hospital or participants' home that followed the principles of a collaborative and evocative motivational interview, favoring the participant's autonomy. The techniques used were open questions, active listening, empathy, returning reflected thoughts, exploring a change in goals, summarizing, and giving feedback. Also, during the enrollment visit, patients were given verbal and written explanation on how to use the app. Free access was granted after receiving an invitation via the hospital health information system (SAP), which prompted the participant to register using an email address as the username. The app could also be downloaded to the carers' phone in case the patient did not have a smartphone.

During the follow-up, the MyPathway app was used by study participants for bidirectional interaction with the research team. It consisted of positive feedback or reinforcement messages in response to the number of hours of NIV use reported by the patient daily. Also, general advice on specific NIV clinical problems was automatically provided by the app according to the patients' weekly input. Additional educational material on physical activity, diet, and sleep hygiene could be accessed at any time via a dedicated link. A web-based clinical portal enabled the research team to monitor the patient-reported NIV hours of use and clinical problems. As indicated, a dedicated nurse with clinical and technical knowledge (one of the authors, MM) took the role of case manager to support collaborative work. She used the web-based portal to identify adherence problems and contacted participants via telephone or at home (for those with severe mobility problems) to enquire about and solve potential clinical or technical problems.

Procedures and Study Outcomes

The primary outcome was a change in self-efficacy, as measured using the Self Efficacy in Sleep Apnea (SEMSA) questionnaire. The SEMSA is a US-designed self-report questionnaire comprised of 26 items that are rated from 1 to 4 on a 4-point Likert scale [30]. The arithmetic mean of the Likert rating for each participant is computed for the overall SEMSA score and each of the 3 factors. The total score ranges from 1 to 4. Higher scores indicate greater risk perception, higher benefit expectancy with treatment, and greater perceived self-efficacy [30].

Secondary outcomes included usability of the ICT tool, as measured using the System Usability Scale [31]; patient satisfaction, as measured using the Net Promoter Score [32] in addition to 3 custom general satisfaction questions measured on a Likert scale; continuity of care, as measured using the Nijmegen Continuity Questionnaire [33]; and the Person-Centred Coordinated Care Experience Questionnaire as described by Leijten et al [34]. Moreover, ventilator-specific data such as the mean hours of daily use, unintentional leaks (L/s), minute ventilation (L/min), tidal volume (mL), and backup rate (breaths/min) were downloaded directly from the NIV machine.

Tertiary outcomes included mortality; health-related quality of life, as measured using the EuroQol 5D questionnaire [35,36]; and sleepiness, as measured using the Epworth Sleepiness Score.

The impact of the motivational mHealth tool recommendations on diet and exercise was indirectly measured by body weight changes.

All assessments were completed at baseline and the final visit scheduled 3 months later. The follow-up was conducted in the outpatient clinic for the control group and remotely by the nurse case manager (MM) using the MyPathway app and its clinical portal for the intervention group. When deemed necessary, the nurse case manager visited the patient at home, or a visit was scheduled in the outpatient clinics. There was no active follow-up for the control group.

Randomization and Masking

All eligible patients were contacted by telephone to briefly explain the study and invite them to participate. Those showing interest were invited to the hospital outpatient clinics. Study investigators (EB, EA, and MM) explained the study face-to-face, and, in case of acceptance, signed consent was obtained. Afterward, the patient was randomized. Before patient enrollment, the randomization scheme was generated using the website randomization.com by one of the researchers (EB). Blocks of 4 were used. Only after the participant provided consent, the investigator opened the envelope with the allocated study group.

Due to the nature of the intervention, neither the participants nor the investigators in direct contact with the participants were blinded. Only the investigator in charge of data analysis was blinded.

Sample Size Calculation, Data Management, and Statistical Analysis

Accepting an α risk of 0.05 and a β risk of 0.2 in a two-sided test, 31 subjects in the intervention group and 31 subjects in the control group were required to achieve a statistically significant difference ≥ 0.35 units in the SEMSA overall score [37]. The common SD was assumed to be 0.46 [38]. A 10% drop-rate was anticipated.

Baseline and end-of-study data (questionnaires) were collected face-to-face at the outpatient clinic by the investigators (EB, EA, and MM). Study data were collected and managed using the REDCap electronic case report form [39,40] hosted at the Hospital Clínic de Barcelona. Data on patient-reported NIV use and clinical problems with NIV were collected online using MyPathway.

Results are presented as mean (SD) or n (%). Comparisons were conducted using Chi-square or Fisher exact tests for categorical variables and Student *t* or Wilcoxon tests, depending on the distribution of the variables, for numerical variables.

Ethics

Study approval was obtained from the Ethics Committee for Clinical Research of Hospital Clínic de Barcelona (HCB/2019/0510). Patients read, understood, and accepted

informed consent, which was signed before enrolment to the study.

Results

Study Population

Between February and March 2019, all patients already being treated with NIV at the noninvasive ventilation clinic at the Hospital Clínic de Barcelona were assessed for eligibility. From an initial sample of 169 eligible patients, 50 (30%) did not meet the inclusion criteria, including 32 who did not have a smartphone or tablet, and 23 (14%) declined participation. Therefore, 67 patients were randomized between February and May 2019 (see the CONSORT flow diagram in [Multimedia Appendix 2](#)). One patient from the intervention group withdrew consent during the trial due to the worsening of his clinical

condition. Baseline demographic and clinical characteristics are shown in [Table 1](#) and [Multimedia Appendix 3](#).

Patient-Reported Outcomes

For the primary outcome, the mean SEMSA score for self-efficacy was not significantly different in the intervention group after the intervention (before: 3.4, SD 0.6; after: 3.4, SD 0.5, $P=.51$).

The perceived risks, outcome expectancies, Epworth Sleepiness Score, and EuroQol 5Q-5D questionnaire score were also not significantly different in the intervention group after the intervention (see [Multimedia Appendix 3](#)). As for the patient experience questionnaires, neither the Nijmegen Continuity Questionnaire nor the Person-Centred Coordinated Care Experience Questionnaire were statistically significantly different between the groups (see [Multimedia Appendix 3](#)).

Table 1. Baseline characteristics of the study groups

	Intervention (n=33)	Control (n=34)	P value
Age (years), mean (SD)	68 (15.8)	65 (14.7)	.31
Male gender, n (%)	19 (58)	19 (58)	>.99
Weight, mean (SD)	86 (31.6)	78 (22.4)	.15
Educational level (n, %)			.73
No schooling	3 (9)	1 (3)	
School education	12 (36)	13 (38)	
Professional formation	17 (52)	19 (56)	
Doctorate or equivalent	1 (3)	1 (3)	
BMI (kg/m ²), mean (SD)	30.5 (7.1)	28.9 (7.4)	.35
Smoking status, n (%)			<.001
Never	12 (36)	16 (49)	
Former	18 (55)	16 (48)	
Current	2 (6)	1 (3)	
Smoking (packs/year), mean (SD)	55.5 (35.7)	52.5 (33)	.003
Diagnostic group, n (%)			
Neuromuscular	4 (12)	8 (24)	.25
Chest wall	11 (33)	10 (30)	.81
Obesity-hypoventilation	5 (15)	5 (15)	>.99
Airway obstructive disease	3 (9)	2 (6)	.66
OSA ^a to CSA ^b	10 (30)	8 (24)	.60
Number of comorbidities per patient, mean (SD)	2 (1.5)	1.8 (1.6)	.68
Comorbidities, %			
Cancer	3	3	>.99
Congestive heart disease	33	27	.60
Ischemic heart disease	24	15	.37
Diabetes	27	36	.47
Stroke	9	9	>.99
Hypertension	67	52	.20
Dementia	3	0	.32
Neurological disorders other than stroke	3	0	.32
Depression/anxiety	18	18	>.99
Dyslipidemia	15	27	.54
Time on noninvasive ventilation (years), mean (SD)	6.75 (6.5)	4.5 (3.5)	.08
AHI ^c , mean (SD)	46 (28.8)	35 (31.6)	.37
CT90 ^d (%), mean (SD)	47 (37.3)	44 (40.4)	.91
Mean ventilatory parameters, mean (SD)			
IPAP ^e (cm H ₂ O)	16 (4.7)	14 (4.7)	.06
EPAP ^f (cm H ₂ O)	7 (2.8)	6 (2.1)	.31
Leak (L/s)	0.05 (0.2)	0.5 (0.09)	.03
Number of hours used per day	7.4 (2)	6.8 (3)	.28

^aOSA: obstructive sleep apnea.

^bCSA: central sleep apnea.

^cAHI: global apnea-hypopnea index for all diagnostic groups.

^dCT90: cumulative sleep time percentage with oxyhemoglobin saturation <90%.

^eIPAP: inspiratory positive airway pressure.

^fEPAP: expiratory positive airway pressure.

Clinical Outcomes

Adherence was measured as the number of hours the NIV was used per day, as recorded by the ventilator. The mean adherence value was not significantly different in the intervention group after the intervention (before: 7.4 hours, SD 2 hours; after: 7.7 hours, SD 2 hours). Mean minute ventilation was the only significantly different ventilatory parameter after the 3-month intervention in the intervention group (before: 7.0 L/min, SD 2 L/min; after: 6.4 L/min, SD 2.1 L/min, $P=.03$). The remaining ventilatory parameters and weight are shown in [Multimedia Appendix 3](#). None of the patients died during the trial.

mHealth Tool Use, Usability, and Acceptability

The Net Promoter Score was -3 (10/33, 31% promoters; 11/33, 34% passives; 11/33, 34% detractors). The 3 Likert-scale questions about the general satisfaction with the app that were rated from 1 (very bad) to 10 (very good) resulted in a mean score of 7.5/10 for the general impression of the app, mean score of 8.2/10 for the user friendliness, and mean score of 8.5/10 for usability of the app without assistance. The mean System Usability Scale score was 78, a reasonably good grading. Up to 42% of the participants used the link to the educational material, and only 18% (6/33) consulted the terms of use. The

mean number of hours of NIV use per day, reported using the mHealth tool, was 7.23 hours (SD 2.48 hours). Use of NIV for more than 4 hours per day during two-thirds of the study period was reported by 45% (15/33) of the patients. Likewise, the reported mean number of days during which NIV was used more than 4 hours in the entire intervention group was 35.6 days (SD 23.6 days). At the end of the study period, 3 participants stopped reporting due to app problems, 1 participant stopped using the app due to health problems, another participant stopped using the app for unknown reasons, and 3 participants decided to use the app on an alternative day basis.

Also, 30% (10/33) of the participants used the app through a family member or carer. It is of note that the nurse case manager was able to solve two-thirds of the technical problems that arose during the first 3 weeks of the study.

The qualitative analysis of the motivational interview as well as the detailed description of the requirements for mHealth to support collaborative work among stakeholders will be reported elsewhere. However, [Table 2](#) summarizes a list of features that the research team agreed were key functional requirements of mHealth tools to effectively support collaborative work among stakeholders involved in home-based respiratory therapies.

Table 2. Requirements to support collaborative work within the noninvasive ventilation service.

Feature	Description of the requirement(s)
Adaptive case management	Capacity to enable the case manager to combine predesigned tasks and approach new cases by reusing structured experiences with previous cases. Over time, the case manager, or other authorized health professionals, should be able to adapt the work plan in a timely fashion to specific patient's requirements without any direct technological support
Team collaboration	Cloud-based, General Data Protection Regulation-compliant, enterprise-proven team collaboration tools to allow patients and health care professionals to break down silos and collaborate seamlessly from any device (mobile phone, tablet, or desktop) towards the health continuum care pathway
Multimedia communication	Enterprise-grade, scalable, high-quality, real-time communication among concurrent participants for file sharing, voice, video, and screen-share sessions with industry-standard encryption
Intelligent bots	Capacity to develop and integrate intelligent bots to guide professionals through continuum care pathways and to improve health risk assessment and service selection
Integration with hospital information systems	Use of HL7 Fast Healthcare Interoperability Resource interoperable middleware to integrate with provider-specific hospital information systems

Discussion

Principal Findings for Patient-Reported Outcomes

We report the results of a behavioral mHealth intervention based on a face-to-face interview and the use of an mHealth tool (MyPathway app) during a 3-month follow-up period with patients with hypercapnic chronic respiratory failure under home-based long-term NIV. To the best of our knowledge, this

is the first randomized controlled trial using digital tools to support behavioral changes in this population [41-44].

In this study, the mean self-efficacy score was already high at baseline ([Table 1](#)), and we did not find a significant effect of the intervention on behavioral changes. Several explanations can be proposed for these results. First, the intervention may need to be more intensive (ie, more than one face-to-face session) [45]. Second, all the participating patients were long-term users without significant sleep symptoms at the time

of enrollment (average use >6 years with an average Epworth Sleepiness Score <10). Therefore, we could hypothesize that behavioral changes had occurred previously, as evidenced by the good average use of NIV (7.4 h/day) and high scores for self-efficacy at baseline. The inclusion of patients who have been newly prescribed NIV in future studies may show a positive impact of the intervention. Third, we may argue that, although NIV use was good among this sample of long-term users, adherence was more a function of necessity or imposition (by family or physicians) than a real feeling of self-management and that most of these chronic patients had not considered initiating behavioral changes [46,47]. Along this line of thought, the population we studied had mobility problems or poor general health, creating barriers for behavioral change [20]. Therefore, any intervention at this stage is likely to be ineffective. This may also be reflected by the lack of interest in consulting the educational material in the app (<50% of the patients did so). Last, we should note that the control group consisted of more patients with neuromuscular pathophysiology. However, the pathophysiology should not affect or have a direct relationship with the measured behavioral outcomes or the capacity and readiness to use the app. Accordingly, educational level is a more important factor [48,49], and both study groups had similar educational levels.

Usability, Acceptability, and Requirements for Supporting Collaborative Work

Notwithstanding the clinical results, it is important to note that the mHealth tool was well received by the patients and their family/caregivers. Despite their complex conditions (2 comorbidities on average) with considerable needs and burdensome treatment, all patients used the app regularly, grading it as generally good, user-friendly, and easy to use without help. Moreover, the System Usability Scale score was good.

As stated in the methods section, we want to highlight the fact that one of the authors (MM) undertook a new professional role during the study period. She became the clinical case manager with additional technical knowledge on the mHealth tool. Patients appreciated this new role very much despite the use of telephone or Whatsapp for bilateral communication. We found that the app lacked this function, and based on our experience, this should become an integral part of any app that includes case management with technical skills. This type of communication functionality should be cloud-based and General Data Protection Regulation-compliant. Moreover, future developments should consider adaptive case management functionality. Also, this communication should be supported by artificial intelligence to help guide professionals through continuum care pathways and improve health risk assessment and service selection. Finally, integration with hospital information systems may facilitate the whole process. This is in line with a recent report on the digital transformation of health care in Europe, which draws upon the experiences of 17 integrated care programs where the importance of communication technologies, new professional roles, and the relevance of clinical workflow evaluation were highlighted [50].

In this respect, we measured 2 process outcomes [51] related to patient experience [52]: continuity of care and person-centered care. Our study population, which included patients as well as their family and carers for one-third of the cases in the intervention group, evaluated both parameters very well. The importance of well-designed clinical workflows with embedded digital health tools may have an impact on not only an NIV service but also other respiratory services. Commonalities include high-complexity patients with clinical and social needs from different stakeholders (eg, physicians, providers, technicians, social workers) and health care tiers (eg, primary care, specialized care). Hernandez et al [24] showed how this complexity can hamper the effectiveness of long-term oxygen therapy. As mentioned, Table 2 shows the proposed elements to overcome the barriers for the successful implementation of digital health tools within clinical workflows relating to respiratory therapies.

Finally, stakeholders play an important role in the design and evaluation of digital health tools [53,54] and, as such, their input should be taken into account when evaluating a service in which there is considerable interplay between patients, different health care tiers, and social and technical services [55]. For an mHealth tool to produce health care value, it should be embedded in the clinical pathways of a well-evaluated clinical service and not as a standalone tool [56].

Strengths and Limitations of the Study

Our study considered the whole population of patients attending the clinic, resulting in a realistic clinical scenario. Another important strength of our study is its potential to demonstrate the positive interaction and collaborative work among the nurse case manager, patients, and family members or caregivers of complex patients using digital health tools. Previous studies [57-59] reported the use of digital tools by family caregivers, emphasizing the importance of including this group of stakeholders, not only as users but also in the co-design process. This stakeholder involvement is also a further step in scaling up digital health tools within clinical workflows [59], which, in our case, were evaluated well. An interesting aspect of our study was the collateral use of qualitative data collected from the motivational interviews and by the nurse case manager during follow-up. The qualitative results presented in Table 2 can be used to support the implementation of mHealth tools in different contexts, keeping in mind the inherent limitations of qualitative research data. We do acknowledge that, by using an existing app, the co-design phase was skipped. Also, we did not measure the technological literacy of our older population (average age 69 years), but, according to Martinez-Alcala et al [60], adults older than 60 years, if highly motivated, are capable of learning and acquiring digital literacy skills. Nonetheless, for some of our older patients (24% were 70-79 years old), especially those with physical limitations (eg, visual impairment), the motivation to learn and exploit all the app functionality was low, although the perceived usefulness was high. This agrees with other reports on the use of technology by older adults [61,62]. Another potential limitation was the heterogeneity of the study population, which directly influenced the mean number of hours of use of the NIV machines and precludes any interpretation. Nonetheless, we observe a strength

in terms of the generalizability of the mHealth tools within the heterogeneous population. Finally, a clear limitation of our study was the exclusion of new NIV patients, where the behavioral intervention may have had more impact. This warrants further study.

Conclusions

The behavioral mHealth intervention explored in this study did not show any effect on self-efficacy, adherence with NIV, or

quality of life in our population of experienced NIV users. Nonetheless, we showed the potential of the mHealth app to manage complex patients and foster collaborative work among stakeholders. Regarding a clinical service that was graded well in terms of continuity of care and person-centered care, in which the needs of relevant stakeholders are properly addressed, we see the potential to further study mHealth tools to induce behavioral change in home-based ventilated patients as well as in other respiratory therapies.

Acknowledgments

We gratefully acknowledge the administrative help from Ms Gemma Guerra. We also want to thank all the patients that participated in this study and to the European Regional Development Fund, FEDER (NEXTCARE project COMRDI15, NextHealth RIS3Cat Community), the FIS-PITES (PI18/00841 and PI17/01068), and the Generalitat de Catalunya (CERCA programme and 2017 SGR-617 specific grant). This work was supported by the European Union's Horizon 2020 Research and Innovation Programme under grant agreement n° GA-689802 (CONNECARE).

Conflicts of Interest

JK and JE are employed by Advanced Digital Innovation (UK) Ltd, the creator of the MyPathway app. The remaining authors have no conflicts of interest to declare.

Multimedia Appendix 1

MyPathway adaptation for home-based non-invasive ventilation.

[\[DOCX File, 562 KB-Multimedia Appendix 1\]](#)

Multimedia Appendix 2

Consort flow diagram.

[\[PDF File \(Adobe PDF File\), 53 KB-Multimedia Appendix 2\]](#)

Multimedia Appendix 3

Baseline characteristics and clinical outcomes tables.

[\[DOCX File, 36 KB-Multimedia Appendix 3\]](#)

Multimedia Appendix 4

CONSORT-eHEALTH checklist (V 1.6.2).

[\[PDF File \(Adobe PDF File\), 101 KB-Multimedia Appendix 4\]](#)

References

1. Lassen H. A PRELIMINARY REPORT ON THE 1952 EPIDEMIC OF POLIOMYELITIS IN COPENHAGEN WITH SPECIAL REFERENCE TO THE TREATMENT OF ACUTE RESPIRATORY INSUFFICIENCY. *The Lancet* 1953 Jan 03;261(6749):37-41. [doi: [10.1016/s0140-6736\(53\)92530-6](https://doi.org/10.1016/s0140-6736(53)92530-6)] [Medline: [13011944](https://pubmed.ncbi.nlm.nih.gov/13011944/)]
2. Clini E, Vitacca M, Foglio K, Simoni P, Ambrosino N. Long-term home care programmes may reduce hospital admissions in COPD with chronic hypercapnia. *Eur Respir J* 1996 Aug;9(8):1605-1610 [FREE Full text] [doi: [10.1183/09031936.96.09081605](https://doi.org/10.1183/09031936.96.09081605)] [Medline: [8866580](https://pubmed.ncbi.nlm.nih.gov/8866580/)]
3. Köhnelein T, Windisch W, Köhler D, Drabik A, Geiseler J, Hartl S, et al. Non-invasive positive pressure ventilation for the treatment of severe stable chronic obstructive pulmonary disease: a prospective, multicentre, randomised, controlled clinical trial. *Lancet Respir Med* 2014 Sep;2(9):698-705. [doi: [10.1016/S2213-2600\(14\)70153-5](https://doi.org/10.1016/S2213-2600(14)70153-5)] [Medline: [25066329](https://pubmed.ncbi.nlm.nih.gov/25066329/)]
4. Bourke SC, Tomlinson M, Williams TL, Bullock RE, Shaw PJ, Gibson GJ. Effects of non-invasive ventilation on survival and quality of life in patients with amyotrophic lateral sclerosis: a randomised controlled trial. *The Lancet Neurology* 2006 Feb;5(2):140-147. [doi: [10.1016/s1474-4422\(05\)70326-4](https://doi.org/10.1016/s1474-4422(05)70326-4)]
5. de Lucas-Ramos P, de Miguel-Díez J, Santacruz-Siminiani A, González-Moro JMR, Buendía-García MJ, Izquierdo-Alonso JL. Benefits at 1 year of nocturnal intermittent positive pressure ventilation in patients with obesity-hypoventilation syndrome. *Respir Med* 2004 Oct;98(10):961-967 [FREE Full text] [doi: [10.1016/j.rmed.2004.03.014](https://doi.org/10.1016/j.rmed.2004.03.014)] [Medline: [15481272](https://pubmed.ncbi.nlm.nih.gov/15481272/)]

6. Budweiser S, Heinemann F, Fischer W, Dobroschke J, Wild PJ, Pfeifer M. Impact of ventilation parameters and duration of ventilator use on non-invasive home ventilation in restrictive thoracic disorders. *Respiration* 2006;73(4):488-494. [doi: [10.1159/000088712](https://doi.org/10.1159/000088712)] [Medline: [16205051](https://pubmed.ncbi.nlm.nih.gov/16205051/)]
7. Ward S, Chatwin M, Heather S, Simonds AK. Randomised controlled trial of non-invasive ventilation (NIV) for nocturnal hypoventilation in neuromuscular and chest wall disease patients with daytime normocapnia. *Thorax* 2005 Dec;60(12):1019-1024 [FREE Full text] [doi: [10.1136/thx.2004.037424](https://doi.org/10.1136/thx.2004.037424)] [Medline: [16299118](https://pubmed.ncbi.nlm.nih.gov/16299118/)]
8. Toussaint M, Chatwin M, Soudon P. Mechanical ventilation in Duchenne patients with chronic respiratory insufficiency: clinical implications of 20 years published experience. *Chron Respir Dis* 2007;4(3):167-177. [doi: [10.1177/1479972307080697](https://doi.org/10.1177/1479972307080697)] [Medline: [17711917](https://pubmed.ncbi.nlm.nih.gov/17711917/)]
9. Lloyd-Owen SJ, Donaldson GC, Ambrosino N, Escarabill J, Farre R, Fauroux B, et al. Patterns of home mechanical ventilation use in Europe: results from the Eurovent survey. *Eur Respir J* 2005 Jun;25(6):1025-1031 [FREE Full text] [doi: [10.1183/09031936.05.00066704](https://doi.org/10.1183/09031936.05.00066704)] [Medline: [15929957](https://pubmed.ncbi.nlm.nih.gov/15929957/)]
10. Laub M, Berg S, Midgren B, Swedish Society of Chest Medicine. Home mechanical ventilation in Sweden--inequalities within a homogenous health care system. *Respir Med* 2004 Jan;98(1):38-42 [FREE Full text] [doi: [10.1016/j.rmed.2003.08.005](https://doi.org/10.1016/j.rmed.2003.08.005)] [Medline: [14959812](https://pubmed.ncbi.nlm.nih.gov/14959812/)]
11. de Lucas Ramos P, Rodríguez González-Moro JM, Paz González L, Santa-Cruz Siminiani A, Cubillo Marcos JM. [Current status of home mechanical ventilation in Spain: results of a national survey]. *Arch Bronconeumol* 2000 Nov;36(10):545-550 [FREE Full text] [Medline: [11149196](https://pubmed.ncbi.nlm.nih.gov/11149196/)]
12. Nonoyama ML, McKim DA, Road J, Guerriere D, Coyte PC, Wasilewski M, et al. Healthcare utilisation and costs of home mechanical ventilation. *Thorax* 2018 Jan 26. [doi: [10.1136/thoraxjnl-2017-211138](https://doi.org/10.1136/thoraxjnl-2017-211138)] [Medline: [29374088](https://pubmed.ncbi.nlm.nih.gov/29374088/)]
13. Mansell SK, Cutts S, Hackney I, Wood MJ, Hawksworth K, Creer DD, et al. Using domiciliary non-invasive ventilator data downloads to inform clinical decision-making to optimise ventilation delivery and patient compliance. *BMJ Open Respir Res* 2018;5(1):e000238 [FREE Full text] [doi: [10.1136/bmjresp-2017-000238](https://doi.org/10.1136/bmjresp-2017-000238)] [Medline: [29531743](https://pubmed.ncbi.nlm.nih.gov/29531743/)]
14. Janssens J, Borel J, Pépin JL, SomnoNIV Group. Nocturnal monitoring of home non-invasive ventilation: the contribution of simple tools such as pulse oximetry, capnography, built-in ventilator software and autonomic markers of sleep fragmentation. *Thorax* 2011 May;66(5):438-445. [doi: [10.1136/thx.2010.139782](https://doi.org/10.1136/thx.2010.139782)] [Medline: [20971980](https://pubmed.ncbi.nlm.nih.gov/20971980/)]
15. Barberan-Garcia A, Ubré M, Roca J, Lacy AM, Burgos F, Risco R, et al. Personalised Prehabilitation in High-risk Patients Undergoing Elective Major Abdominal Surgery: A Randomized Blinded Controlled Trial. *Ann Surg* 2018 Jan;267(1):50-56. [doi: [10.1097/SLA.0000000000002293](https://doi.org/10.1097/SLA.0000000000002293)] [Medline: [28489682](https://pubmed.ncbi.nlm.nih.gov/28489682/)]
16. Troosters T, Maltais F, Leidy N, Lavoie KL, Sedeno M, Janssens W, et al. Effect of Bronchodilation, Exercise Training, and Behavior Modification on Symptoms and Physical Activity in Chronic Obstructive Pulmonary Disease. *Am J Respir Crit Care Med* 2018 Oct 15;198(8):1021-1032. [doi: [10.1164/rccm.201706-1288OC](https://doi.org/10.1164/rccm.201706-1288OC)] [Medline: [29664681](https://pubmed.ncbi.nlm.nih.gov/29664681/)]
17. Slovinec D'Angelo ME, Pelletier LG, Reid RD, Huta V. The roles of self-efficacy and motivation in the prediction of short- and long-term adherence to exercise among patients with coronary heart disease. *Health Psychol* 2014 Nov;33(11):1344-1353. [doi: [10.1037/hea0000094](https://doi.org/10.1037/hea0000094)] [Medline: [25133848](https://pubmed.ncbi.nlm.nih.gov/25133848/)]
18. Kenyon DB, Kubik MY, Davey C, Sirard J, Fulkerson JA. Alternative high school students' physical activity: role of self-efficacy. *Am J Health Behav* 2012 Mar;36(3):300-310 [FREE Full text] [doi: [10.5993/AJHB.36.3.2](https://doi.org/10.5993/AJHB.36.3.2)] [Medline: [22370432](https://pubmed.ncbi.nlm.nih.gov/22370432/)]
19. Teixeira PJ, Marques MM. Health Behavior Change for Obesity Management. *Obes Facts* 2017;10(6):666-673 [FREE Full text] [doi: [10.1159/000484933](https://doi.org/10.1159/000484933)] [Medline: [29237167](https://pubmed.ncbi.nlm.nih.gov/29237167/)]
20. McAuley E, Szabo A, Gothe N, Olson EA. Self-efficacy: Implications for Physical Activity, Function, and Functional Limitations in Older Adults. *Am J Lifestyle Med* 2011 Jul;5(4) [FREE Full text] [doi: [10.1177/1559827610392704](https://doi.org/10.1177/1559827610392704)] [Medline: [24353482](https://pubmed.ncbi.nlm.nih.gov/24353482/)]
21. Bandura A. Self-efficacy mechanism in human agency. *Am Psychol* 1982;37(2):122-147. [Medline: [15064770](https://pubmed.ncbi.nlm.nih.gov/15064770/)]
22. Bandura A. Self-efficacy: toward a unifying theory of behavioral change. *Psychol Rev* 1977 Mar;84(2):191-215. [doi: [10.1037//0033-295x.84.2.191](https://doi.org/10.1037//0033-295x.84.2.191)] [Medline: [847061](https://pubmed.ncbi.nlm.nih.gov/847061/)]
23. Schwarzer R, Fuchs R. Self-efficacy and health behaviours. In: Conner M, Norman P, editors. *Predicting health behaviour: Research and practice with social cognition models*. Philadelphia: Open University Press; 1996:163-196.
24. Hernandez C, Aibar J, de Batlle J, Gomez-Cabrero D, Soler N, Duran-Tauleria E, NEXES Consortium. Assessment of health status and program performance in patients on long-term oxygen therapy. *Respir Med* 2015 Apr;109(4):500-509 [FREE Full text] [doi: [10.1016/j.rmed.2015.01.005](https://doi.org/10.1016/j.rmed.2015.01.005)] [Medline: [25771036](https://pubmed.ncbi.nlm.nih.gov/25771036/)]
25. Cano I, Dueñas-Espín I, Hernandez C, de Batlle J, Benavent J, Contel JC, et al. Protocol for regional implementation of community-based collaborative management of complex chronic patients. *NPJ Prim Care Respir Med* 2017 Jul 14;27(1):44 [FREE Full text] [doi: [10.1038/s41533-017-0043-9](https://doi.org/10.1038/s41533-017-0043-9)] [Medline: [28710482](https://pubmed.ncbi.nlm.nih.gov/28710482/)]
26. Cano I, Alonso A, Hernandez C, Burgos F, Barberan-Garcia A, Roldan J, et al. An adaptive case management system to support integrated care services: Lessons learned from the NEXES project. *J Biomed Inform* 2015 Jun;55:11-22 [FREE Full text] [doi: [10.1016/j.jbi.2015.02.011](https://doi.org/10.1016/j.jbi.2015.02.011)] [Medline: [25796455](https://pubmed.ncbi.nlm.nih.gov/25796455/)]

27. Ambrosino N, Vitacca M, Dreher M, Isetta V, Montserrat JM, Tonia T, ERS Tele-Monitoring of Ventilator-Dependent Patients Task Force. Tele-monitoring of ventilator-dependent patients: a European Respiratory Society Statement. *Eur Respir J* 2016 Sep;48(3):648-663 [FREE Full text] [doi: [10.1183/13993003.01721-2015](https://doi.org/10.1183/13993003.01721-2015)] [Medline: [27390283](https://pubmed.ncbi.nlm.nih.gov/27390283/)]
28. COMMUNICATION FROM THE COMMISSION TO THE EUROPEAN PARLIAMENT, THE COUNCIL, THE EUROPEAN ECONOMIC AND SOCIAL COMMITTEE AND THE COMMITTEE OF THE REGIONS eHealth Action Plan 2012-2020 - Innovative healthcare for the 21st century /* COM/2012/0736 final */ Internet. OPOCE; URL: <https://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:52012DC0736&from=EN> [accessed 2018-07-31]
29. MyPathway. URL: <https://mypathway.healthcare/> [accessed 2019-03-01]
30. Montplaisir J, Bédard MA, Richer F, Rouleau I. Neurobehavioral manifestations in obstructive sleep apnea syndrome before and after treatment with continuous positive airway pressure. *Sleep* 1992 Dec;15(6 Suppl):S17-S19. [doi: [10.1093/sleep/15.suppl_6.s17](https://doi.org/10.1093/sleep/15.suppl_6.s17)] [Medline: [1470802](https://pubmed.ncbi.nlm.nih.gov/1470802/)]
31. Sauro J. Measuring Usability With The System Usability Scale (SUS). URL: <https://measuringu.com/umux-lite/> [accessed 2019-02-01]
32. Reichheld F. The One Number You Need to Grow. *Harv Bus Rev* 2003;81(12):124 [FREE Full text]
33. Uijen AA, Schellevis FG, van den Bosch WJHM, Mokkink HGA, van Weel C, Schers HJ. Nijmegen Continuity Questionnaire: development and testing of a questionnaire that measures continuity of care. *J Clin Epidemiol* 2011 Dec;64(12):1391-1399. [doi: [10.1016/j.jclinepi.2011.03.006](https://doi.org/10.1016/j.jclinepi.2011.03.006)] [Medline: [21689904](https://pubmed.ncbi.nlm.nih.gov/21689904/)]
34. Leijten FRM, Hoedemakers M, Struckmann V, Kraus M, Cheraghi-Sohi S, Zemplényi A, SELFIE consortium. Defining good health and care from the perspective of persons with multimorbidity: results from a qualitative study of focus groups in eight European countries. *BMJ Open* 2018 Aug 30;8(8):e021072 [FREE Full text] [doi: [10.1136/bmjopen-2017-021072](https://doi.org/10.1136/bmjopen-2017-021072)] [Medline: [30166294](https://pubmed.ncbi.nlm.nih.gov/30166294/)]
35. Wolowacz SE, Briggs A, Belozeroff V, Clarke P, Doward L, Goeree R, et al. Estimating Health-State Utility for Economic Models in Clinical Studies: An ISPOR Good Research Practices Task Force Report. *Value Health* 2016;19(6):704-719 [FREE Full text] [doi: [10.1016/j.jval.2016.06.001](https://doi.org/10.1016/j.jval.2016.06.001)] [Medline: [27712695](https://pubmed.ncbi.nlm.nih.gov/27712695/)]
36. Pickard AS, Wilke C, Jung E, Patel S, Stavem K, Lee TA. Use of a preference-based measure of health (EQ-5D) in COPD and asthma. *Respir Med* 2008 Apr;102(4):519-536 [FREE Full text] [doi: [10.1016/j.rmed.2007.11.016](https://doi.org/10.1016/j.rmed.2007.11.016)] [Medline: [18180151](https://pubmed.ncbi.nlm.nih.gov/18180151/)]
37. Wyrwich KW. Minimal important difference thresholds and the standard error of measurement: is there a connection? *J Biopharm Stat* 2004 Feb;14(1):97-110. [doi: [10.1081/BIP-120028508](https://doi.org/10.1081/BIP-120028508)] [Medline: [15027502](https://pubmed.ncbi.nlm.nih.gov/15027502/)]
38. Philip P, Bioulac S, Altena E, Morin CM, Ghorayeb I, Coste O, et al. Specific insomnia symptoms and self-efficacy explain CPAP compliance in a sample of OSAS patients. *PLoS One* 2018;13(4):e0195343 [FREE Full text] [doi: [10.1371/journal.pone.0195343](https://doi.org/10.1371/journal.pone.0195343)] [Medline: [29617414](https://pubmed.ncbi.nlm.nih.gov/29617414/)]
39. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009 Apr;42(2):377-381 [FREE Full text] [doi: [10.1016/j.jbi.2008.08.010](https://doi.org/10.1016/j.jbi.2008.08.010)] [Medline: [18929686](https://pubmed.ncbi.nlm.nih.gov/18929686/)]
40. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, REDCap Consortium. The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform* 2019 Jul;95:103208. [doi: [10.1016/j.jbi.2019.103208](https://doi.org/10.1016/j.jbi.2019.103208)] [Medline: [31078660](https://pubmed.ncbi.nlm.nih.gov/31078660/)]
41. Hamine S, Gerth-Guyette E, Faulx D, Green B, Ginsburg A. Impact of mHealth chronic disease management on treatment adherence and patient outcomes: a systematic review. *J Med Internet Res* 2015 Feb 24;17(2):e52 [FREE Full text] [doi: [10.2196/jmir.3951](https://doi.org/10.2196/jmir.3951)] [Medline: [25803266](https://pubmed.ncbi.nlm.nih.gov/25803266/)]
42. Whitehead L, Seaton P. The Effectiveness of Self-Management Mobile Phone and Tablet Apps in Long-term Condition Management: A Systematic Review. *J Med Internet Res* 2016 May 16;18(5):e97 [FREE Full text] [doi: [10.2196/jmir.4883](https://doi.org/10.2196/jmir.4883)] [Medline: [27185295](https://pubmed.ncbi.nlm.nih.gov/27185295/)]
43. Lee J, Choi M, Lee S, Jiang N. Effective behavioral intervention strategies using mobile health applications for chronic disease management: a systematic review. *BMC Med Inform Decis Mak* 2018 Feb 20;18(1):12 [FREE Full text] [doi: [10.1186/s12911-018-0591-0](https://doi.org/10.1186/s12911-018-0591-0)] [Medline: [29458358](https://pubmed.ncbi.nlm.nih.gov/29458358/)]
44. Yi JY, Kim Y, Cho Y, Kim H. Self-management of Chronic Conditions Using mHealth Interventions in Korea: A Systematic Review. *Healthc Inform Res* 2018 Jul;24(3):187-197 [FREE Full text] [doi: [10.4258/hir.2018.24.3.187](https://doi.org/10.4258/hir.2018.24.3.187)] [Medline: [30109152](https://pubmed.ncbi.nlm.nih.gov/30109152/)]
45. Voils CI, King HA, Maciejewski ML, Allen KD, Yancy WS, Shaffer JA. Approaches for informing optimal dose of behavioral interventions. *Ann Behav Med* 2014 Dec;48(3):392-401 [FREE Full text] [doi: [10.1007/s12160-014-9618-7](https://doi.org/10.1007/s12160-014-9618-7)] [Medline: [24722964](https://pubmed.ncbi.nlm.nih.gov/24722964/)]
46. Mogler BK, Shu SB, Fox CR, Goldstein NJ, Victor RG, Escarce JJ, et al. Using insights from behavioral economics and social psychology to help patients manage chronic diseases. *J Gen Intern Med* 2013 May;28(5):711-718 [FREE Full text] [doi: [10.1007/s11606-012-2261-8](https://doi.org/10.1007/s11606-012-2261-8)] [Medline: [23229906](https://pubmed.ncbi.nlm.nih.gov/23229906/)]
47. Janz NK, Becker MH. The Health Belief Model: a decade later. *Health Educ Q* 1984;11(1):1-47. [doi: [10.1177/109019818401100101](https://doi.org/10.1177/109019818401100101)] [Medline: [6392204](https://pubmed.ncbi.nlm.nih.gov/6392204/)]
48. Hsu W, Chiang C, Yang S. The effect of individual factors on health behaviors among college students: the mediating effects of eHealth literacy. *J Med Internet Res* 2014 Dec 12;16(12):e287 [FREE Full text] [doi: [10.2196/jmir.3542](https://doi.org/10.2196/jmir.3542)] [Medline: [25499086](https://pubmed.ncbi.nlm.nih.gov/25499086/)]

49. Neter E, Brainin E. eHealth literacy: extending the digital divide to the realm of health information. *J Med Internet Res* 2012 Jan 27;14(1):e19 [FREE Full text] [doi: [10.2196/jmir.1619](https://doi.org/10.2196/jmir.1619)] [Medline: [22357448](https://pubmed.ncbi.nlm.nih.gov/22357448/)]
50. Baltaxe E, Cypionka T, Kraus M, Reiss M, Askildsen JE, Grenkovic R, et al. Digital Health Transformation of Integrated Care in Europe: Overarching Analysis of 17 Integrated Care Programs. *J Med Internet Res* 2019 Sep 26;21(9):e14956 [FREE Full text] [doi: [10.2196/14956](https://doi.org/10.2196/14956)] [Medline: [31573914](https://pubmed.ncbi.nlm.nih.gov/31573914/)]
51. Donabedian A. The Quality of Care. *JAMA* 1988 Sep 23;260(12):1743-1748 [FREE Full text] [doi: [10.1001/jama.1988.03410120089033](https://doi.org/10.1001/jama.1988.03410120089033)]
52. Berwick DM, Nolan TW, Whittington J. The triple aim: care, health, and cost. *Health Aff (Millwood)* 2008;27(3):759-769. [doi: [10.1377/hlthaff.27.3.759](https://doi.org/10.1377/hlthaff.27.3.759)] [Medline: [18474969](https://pubmed.ncbi.nlm.nih.gov/18474969/)]
53. Taylor MJ, McNicholas C, Nicolay C, Darzi A, Bell D, Reed JE. Systematic review of the application of the plan-do-study-act method to improve quality in healthcare. *BMJ Qual Saf* 2014 Apr;23(4):290-298 [FREE Full text] [doi: [10.1136/bmjqs-2013-001862](https://doi.org/10.1136/bmjqs-2013-001862)] [Medline: [24025320](https://pubmed.ncbi.nlm.nih.gov/24025320/)]
54. Ploeg J, Ali MU, Markle-Reid M, Valaitis R, Bartholomew A, Fitzpatrick-Lewis D, et al. Caregiver-Focused, Web-Based Interventions: Systematic Review and Meta-Analysis (Part 2). *J Med Internet Res* 2018 Oct 26;20(10):e11247 [FREE Full text] [doi: [10.2196/11247](https://doi.org/10.2196/11247)] [Medline: [30368439](https://pubmed.ncbi.nlm.nih.gov/30368439/)]
55. Baltaxe E, Cano I, Herranz C, Barberan-Garcia A, Hernandez C, Alonso A, et al. Evaluation of integrated care services in Catalonia: population-based and service-based real-life deployment protocols. *BMC Health Serv Res* 2019 Jun 11;19(1):370 [FREE Full text] [doi: [10.1186/s12913-019-4174-2](https://doi.org/10.1186/s12913-019-4174-2)] [Medline: [31185997](https://pubmed.ncbi.nlm.nih.gov/31185997/)]
56. Lennon MR, Bouamrane M, Devlin AM, O'Connor S, O'Donnell C, Chetty U, et al. Readiness for Delivering Digital Health at Scale: Lessons From a Longitudinal Qualitative Evaluation of a National Digital Health Innovation Program in the United Kingdom. *J Med Internet Res* 2017 Feb 16;19(2):e42 [FREE Full text] [doi: [10.2196/jmir.6900](https://doi.org/10.2196/jmir.6900)] [Medline: [28209558](https://pubmed.ncbi.nlm.nih.gov/28209558/)]
57. Piette JD, Striplin D, Marinec N, Chen J, Trivedi RB, Aron DC, et al. A Mobile Health Intervention Supporting Heart Failure Patients and Their Informal Caregivers: A Randomized Comparative Effectiveness Trial. *J Med Internet Res* 2015 Jun 10;17(6):e142 [FREE Full text] [doi: [10.2196/jmir.4550](https://doi.org/10.2196/jmir.4550)] [Medline: [26063161](https://pubmed.ncbi.nlm.nih.gov/26063161/)]
58. Lapid MI, Atherton PJ, Clark MM, Kung S, Sloan JA, Rummans TA. Cancer Caregiver: Perceived Benefits of Technology. *Telemed J E Health* 2015 Nov;21(11):893-902 [FREE Full text] [doi: [10.1089/tmj.2014.0117](https://doi.org/10.1089/tmj.2014.0117)] [Medline: [26075800](https://pubmed.ncbi.nlm.nih.gov/26075800/)]
59. Costa Stutzel M, Filippo MP, Sztajnberg A, da Costa RMEM, Brites ADS, da Motta LB, et al. Multi-part quality evaluation of a customized mobile application for monitoring elderly patients with functional loss and helping caregivers. *BMC Med Inform Decis Mak* 2019 Jul 22;19(1):140 [FREE Full text] [doi: [10.1186/s12911-019-0839-3](https://doi.org/10.1186/s12911-019-0839-3)] [Medline: [31331309](https://pubmed.ncbi.nlm.nih.gov/31331309/)]
60. Martínez-Alcalá CI, Rosales-Lagarde A, Alonso-Lavernia MDA, Ramírez-Salvador JA, Jiménez-Rodríguez B, Cepeda-Rebollar RM, et al. Digital Inclusion in Older Adults: A Comparison Between Face-to-Face and Blended Digital Literacy Workshops Internet. *Frontiers in ICT* 2018;5:21 [FREE Full text] [doi: [10.3389/fict.2018.00021](https://doi.org/10.3389/fict.2018.00021)]
61. Parker SJ, Jessel S, Richardson JE, Reid MC. Older adults are mobile too! Identifying the barriers and facilitators to older adults' use of mHealth for pain management. *BMC Geriatr* 2013 May 06;13:43 [FREE Full text] [doi: [10.1186/1471-2318-13-43](https://doi.org/10.1186/1471-2318-13-43)] [Medline: [23647949](https://pubmed.ncbi.nlm.nih.gov/23647949/)]
62. Spann A, Stewart E. Barriers and facilitators of older people's mHealth usage: A qualitative review of older people's views. *Human Technology* 2018 Nov 30;14(3):264-296. [doi: [10.17011/ht/urn.201811224834](https://doi.org/10.17011/ht/urn.201811224834)]

Abbreviations

ICT: information and communication technologies

mHealth: mobile health

NIV: noninvasive ventilation

SEMSA: Self Efficacy in Sleep Apnea

Edited by G Eysenbach; submitted 25.09.19; peer-reviewed by W Zhang, J Xu, J Aikens; comments to author 25.11.19; revised version received 19.01.20; accepted 04.02.20; published 13.04.20

Please cite as:

Baltaxe E, Embid C, Aumatell E, Martínez M, Barberan-Garcia A, Kelly J, Eaglesham J, Herranz C, Vargiu E, Montserrat JM, Roca J, Cano I

Integrated Care Intervention Supported by a Mobile Health Tool for Patients Using Noninvasive Ventilation at Home: Randomized Controlled Trial

JMIR Mhealth Uhealth 2020;8(4):e16395

URL: <http://mhealth.jmir.org/2020/4/e16395/>

doi: [10.2196/16395](https://doi.org/10.2196/16395)

PMID:

©Erik Baltaxe, Cristina Embid, Eva Aumatell, María Martínez, Anael Barberan-Garcia, John Kelly, John Eaglesham, Carmen Herranz, Eloisa Vargiu, Josep Maria Montserrat, Josep Roca, Isaac Cano. Originally published in JMIR mHealth and uHealth (<http://mhealth.jmir.org>), 13.04.2020. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR mHealth and uHealth, is properly cited. The complete bibliographic information, a link to the original publication on <http://mhealth.jmir.org/>, as well as this copyright and license information must be included.